arXiv:1606.04623v1 [hep-ph] 15 Jun 2016

$\bar{K}$ -induced formation of the $f_2(1270)$ and $f_2'(1525)$ resonances on proton targets

Yin Huang,[1,2,3,*] Ju-jun Xie,[1,3,†] Jun He,[1,3] Xurong Chen,[1,3] and Hong-fei Zhang[2,3]

[1]*Research Center for Hadron and CSR Physics, Lanzhou University and Institute of Modern Physics of CAS, Lanzhou 730000,China*
[2]*School of Nuclear Science and Technology, Lanzhou University, Lanzhou 730000, China*
[3]*Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou 730000, China*

(Dated: June 16, 2016)

We investigate the productiong of $f_2(1270)$ and $f_2'(1525)$ mesons in the $K^-p \to \Lambda f_2(1270)$,$K^-p \to \Lambda f_2'(1525)$ and $K^-p \to K^+K^-\Lambda$ reactions within an effective Lagrangian approach. For $K^-p \to f_2\Lambda$ reaction, by considering the contributions from the *t*-channel K^+ exchange and *u*-channel nucleon pole, we get a fairly good description of the experimental measurements about the total and differential cross sections. Based on the studies of the $K^-p \to f_2\Lambda$ reaction, we investigate $K^-p \to K^+K^-\Lambda$ reaction including the contributions from the $f_2(1270)$ and $f_2'(1525)$ mesons decaying into K^+K^- pair. The total cross sections and invariant mass distribution of the $Kp \to K^+K^-\Lambda$ reaction are predicted. The results can be tested in future experiments and therefore offer new clues on the nature of these tensor states.

PACS numbers: 13.60.Le, 12.39.Mk,13.25.Jx

I. INTRODUCTION

Since the charmonium-like resonance *XYZ* and the new baryonic P_c [1] states are observed, a lot of experiments spring up to study the exotic states from Belle, BaBar, BESIII, LHCb, CDF, D0 and other collaborations. And people believe that the traditional convention, the meson is made up of quark and antiquark as well as baryon is made up of three quarks, is broken. Furthermore, it is not surprising to find out that many low-lying states, even those long believed to be conventional $q\bar{q}$ (or qqq) states, may have large components of other nature. These observation has attracted a lot of attention from the theoretical side. Various explanations of these states have been proposed, such as molecules, mutiquark states, kinematic effects, or mixtures of components of different nature. Nevertheless, up to now none of them has been accepted unanimously.

Indeed, it has been shown that many of the lowlying mesonic states can be understood not only as $q\bar{q}$ states but also as meson-meson molecules, dynamically generated in the so-called unitary approaches. For example, $f_2(1270)$, $f_2'(1525)$ and $K_2^*(1430)$ are listed in the Particle Data Group [2] as the lightest tensor states and correspond to the 2^{++} ground-state nonet. On the other hand, these states have been studied tensor mesons in the unitary approach based on the hidden-gauge Lagrangians. It was found that the $f_2(1270)$, $f'(1525)$, and $K_2^*(1430)$ are dynamically generated from vector-meson-vector-meson interactions [3]. Within this picture, the branching ratios into pseudoscalar-pseudoscalar and vector-vector final states are all consistent with data [3]. In Ref. [4], the two-photon decay widths of the $f_0(1370)$ and $f_2(1270)$ have been calculated and found to agree with data. Furthermore, the ratios of the J/ψ decay rates into a vector meson (ϕ, ω,or K^*) and one of the tensor states [$f_2(1270)$, $f_2'(1525)$, and $K_2^*(1430)$] have been calculated, and the agreement with data is found to be quite reasonable [5]. Following the same approach, in Ref. [6] it is shown that the ratio of the J/ψ decay rates into $\gamma f_2(1270)$ and $\gamma f_2'(1525)$ also agrees with data. The agreement with experimental data turns out to be quite good in general, providing support to the underlying assumption that these states contain large meson-meson components.

Assuming that the $f_2(1270)$ resonance is a $\rho-\rho$ molecular state, it was found in Ref. [7] that the differential cross section as well as the t dependence of the $\gamma p \to p f_2(1270)$ reaction are in good agreement with the experimental results [8] and provide support for the molecular picture of the $f_2(1270)$ resonance in the first baryonic reaction. In a recent work [9], taking the $f_2'(1525)$ resonances are dynamically generated states from vector-meson-vector-meson interactions in the *s*-wave with spin $S=2$, the $\gamma p \to f_2'(1525)p$ reaction has been studied.

In addition to the photoproduction processes, the $\bar{K}N$ scattering can also provide important information on tensor mesons. Hence, we study the $Kp \to \Lambda f_2$ ($f_2 \equiv f_2(1270)$, $f_2'(1525)$) reaction within an effective Lagrangian approach in this work. The contributions from the Born terms including $t-$channel K^+ exchange and *u*-channel nucleon pole are considered. Furthermore, for the low energy of the $K^-p \to K^+K^-\Lambda$ reaction, we pay especially attention on the role of the $f_2(1270)$ and $f_2'((1525)$ mesons.

This paper is organized as follows. After the introduction, we present the calculation of the production of $f_2(1270)$, $f_2'(1525)$ via $\bar{K}$ -induced formation on proton targets. In Sec. III, the contribution to the $K^+K^-\Lambda$ final state is discussed, and the corresponding total cross sections and invariant mass distribution of these reactions are given. This work ends with the conclusion.

II. THE $K^-p \to f_2(1270)\Lambda$, $f_2'(1525)\Lambda$ REACTION

We choose the production process $K^-p \to \Lambda f_2$ reactions to study the production of $f_2(1270)$ and $f_2'(1525)$ which can couple to K^+K^- [2]. First, we mainly concentrate on the production probability of f_2 in the $K^-p \to f_2\Lambda$ process, where

*Electronic address: huangy2014@lzu.cn
†Electronic address: xiejujun@impcas.ac.cn



FIG. 1: Feynman diagrams for $K^- p \to f_2(1270)\Lambda$, $f_2'(1525)\Lambda$ reaction. The contributions from t-channel K^+ exchange (a), u-channel nucleon pole (b). we also show the definition of the kinematical (p_1, p_2, p_3, q, p_5) that we use in the present calculation. In addition, we use $k_t = p_1 - q$, and $k_u = p_1 - p_5$.



FIG. 2: (color online). The obtained total cross section for $K^- p \to f_2(1270)\Lambda$ with the typical cutoff $\Lambda_K = 0.51 - 0.53$ GeV. The experimental data are taken from Ref. [10](circle) and Ref. [11](square)

the total cross sections of these process are discussed.

The basic tree level Feynman diagrams for the $K^- p \to f_2\Lambda$ reaction are shown in Fig. 1, where the contributions from the



FIG. 3: The obtained total cross section for $K^- p \to f_2'(1525)\Lambda$ with the typical cutoff $\Lambda_K = 0.51 - 0.53$ GeV. The experimental data are taken from Ref. [10](circle),Ref. [11](star) and Ref. [12](square)

$t-$channel K^+ exchange process [Fig. 1(a)], and the nucleon pole [Fig. 1(b)] are taken into account. While the s-channel processes are neglected since the information of those processes is scarce and we expect these contributions to be small. To compute the amplitudes of these diagrams shown in Fig. 1, the effective Lagrangian densities for the relevant interaction vertexes are needed. We adopt the effective Lagrangians as used in Refs. [13–16],

$$\mathcal{L}_{Kp\Lambda} = \frac{g_{Kp\Lambda}}{M_N + M_\Lambda}\bar{p}\gamma^\mu\gamma_5\Lambda\partial_\mu K^+ + H.c., \quad (1)$$

$$\mathcal{L}_{f_2KK} = \frac{g_{f_2K^+K^-}}{2m_{f_2}} f_{2\mu\nu}[K^-\partial_\mu\partial_\nu K^+ + \partial_\mu\partial_\nu K^- K^+ - 2\partial_\mu K^+\partial_\nu K^-] + H.c.,$$

$$\mathcal{L}_{f_2NN} = i\frac{G_{f_2NN}}{2M_N}\bar{N}(\gamma_\mu\partial_\nu + \gamma_\nu\partial_\mu)N f_2^{\mu\nu} + \frac{F_{f_2NN}}{(2M_N)^2}\partial_\mu\bar{N}\partial_\nu N f_2^{\mu\nu} + H.c., \quad (2)$$

where M_N and M_Λ are the masses of the nucleon and $\Lambda(1115)$, respectively. The $p, \Lambda, f_{2\mu\nu}$ are the nucleon field, $\Lambda(1115)$ filed, and f_2 filed, respectively. The K is kaon field and the isodoublets are defined as

$$K = \begin{pmatrix} K^+ \\ K^0 \end{pmatrix}; \bar{K} = \begin{pmatrix} K^-, \bar{K}^0 \end{pmatrix}. \quad (3)$$

The coupling constant $g_{Kp\Lambda}$ can be determined by flavor $SU(3)$ symmetry relations, which give $g_{Kp\Lambda} = 13.24$ [13]. While the value of the coupling constant $g_{f_2K^+K^-}$ can be determined from the partial decay width of $f_2 \to K^+K^-$, which can be obtained from Eq. (2),

$$\Gamma_{f_2\to K^-K^+} = \frac{m_{f_2}}{480\pi} g^2_{f_2K^+K^-}(1 - \frac{4m_K^2}{m_{f_2}^2})^{5/2}, \quad (4)$$



FIG. 4: The obtained differential cross section for $K^- p \to f_2(1270)\Lambda$ with the typical cutoff $\Lambda_K = 0.51 - 0.53$ GeV. The experimental data denoted by the squares are taken from Ref. [11].

TABLE I: The coupling constants g_{f_2KK}, G_{f_2NN},and F_{f_2NN} for $f_2 \to K\bar{K}$ and $N\bar{N}$ reaction.

Resonance	*channel*	g_{f_2KK}	G_{f_2NN}	F_{f_2NN}
$f_2(1270)$	$K\bar{K}$	9.96	−−	−−
	$N\bar{N}$	−−	2.19	0
$f_2'(1525)$	$K\bar{K}$	15.78	−−	−−
	$N\bar{N}$	−−	0	0

where m_{f_2} and m_K are mass of f_2 and kaon, respectively.

With mass (m_{f_2} = 1.275 GeV, 1.525 GeV, m_K = 0.494 GeV), total decay width ($\Gamma_{f_2(1270)}$ = 0.185 GeV, $\Gamma_{f_2(1525)}$ = 0.073 GeV), and decay branching ratio of $f_2 \to K^+K^-$ [Br($f_2(1270) \to K^+K^-$) = 0.046, Br($f_2(1525) \to K^+K^-$) = 0.887], from Eq. (4), we obtain the coupling constant are shown in Tab. I.

By using the tensor meson dominance (TMD) [17], one can determine the universal coupling constant of the f_2 meson from its decay into two pions, which can then be used to determine the coupling constant for $f_2 \to N\bar{N}$ reaction (See Ref. [18] for details). And are shown in Table. I.



FIG. 5: The obtained differential cross section for $K^- p \to f_2'(1525)\Lambda$ with the typical cutoff $\Lambda_K = 0.51 - 0.53$ GeV. The experimental data are taken from Ref. [11].

In evaluating the scattering amplitudes of the $K^- p \to f_2\Lambda$ reaction, we need to include the form factors because the hadrons are not pointlike particles. For the t-channel K^+ meson exchange, we adopt here a common scheme used in many previous works [19]

$$F_K(k_t^2) = \frac{\Lambda_K^2 - m_k^2}{\Lambda_K^2 - k_t^2}, \tag{5}$$

where Λ_K is the cutoff parameter.

For the s-channel and u-channel processes, we adopt a form factor [19],

$$\mathcal{F}_B(q_{ex}^2, M_{ex}) = \frac{\Lambda_B^4}{\Lambda^4 + (q_{ex}^2 - M_{ex}^2)^2} \tag{6}$$

where q_{ex} and M_{ex} are the four-momentum and the mass of the exchanged hadron, respectively. The cut-off parameter is constrained between 0.8 and 1.5 GeV for all channels. For simplicity, we take Λ_B = 0.98 GeV [19] for the u-channel nucleon pole, and the s-channel resonance exchanges.

With the above preparation, we finally obtain the amplitude of the $K^-(p_1)p(p_2) \to f_2(q)\Lambda(p_5)$ process,

$$\mathcal{T}_{fi}^a = \frac{-i}{2m_{f_2}} \frac{g_{Kp\Lambda} g_{f_2K^+K^-}}{M_N + M_\Lambda} \bar{u}(p_5, s_5) \not{k}_t \gamma_5 u(p_2, s_2) \frac{1}{k_t^2 - m_K^2} F_K(k_t^2) \times T_{f_2}^{*\mu\nu}(q, \lambda)[-k_{t\mu}k_{t\nu} - p_{1\mu}p_{1\nu} + 2k_{t\mu}p_{1\nu}]. \tag{7}$$

$$\mathcal{T}_{fi}^b = -i \frac{g_{Kp\Lambda} G_{f_{2NN}}}{2M_N(M_N + M_\Lambda)} \bar{u}(p_5, s_5) \not{p}_1 \gamma_5 \frac{\not{k}_u + M_N}{k_u^2 - M_N^2} \times (\gamma_\mu p_{2\nu} + \gamma_\nu p_{2\mu}) T_{f_2}^{*\mu\nu}(q, \lambda) u(p_2, s_2) \mathcal{F}_N(k_u^2, M_N), \tag{8}$$

where s_5, p_5 and s_2, p_2 denote the spin polarization variables and the four-momenta of the outgoing Λ and the initial proton, respectively, while q, λ are the four-momenta and spin polarization variables of the f_2 meson. The $\bar{u}(p_5, s_5)$ and $u(p_2, s_2)$

are the Dirac spinors for the Λ and proton, respectively, while the $T^{*\mu\nu}_{f_2}(q,\lambda)$ is the polarization tensor of the f_2.

By defining $t=(p_1-q)^2=k_t^2, s=(p_1+p_2)^2$, the corresponding unpolarized differential cross section reads as

$$\frac{d\sigma}{dt}=\frac{M_N M_\Lambda}{16\pi s}\frac{1}{|\vec{p}_{1cm}|^2}(\frac{1}{2}\sum_{s_2,s_5,\lambda}|\mathcal{T}_{fi}|^2). \quad (9)$$

The total cross section can be obtained by integrating over the range of $|\vec{k}_t|$.

The sum over polarizations, in Eq. (4), can be easily done thanks to

$$\sum_\lambda T_{\mu\nu}(q,\lambda)T^*_{\alpha\beta}(q,\lambda)=P_{\mu\nu\alpha\beta}$$
$$=\frac{1}{2}(\tilde{g}_{\mu\alpha}\tilde{g}_{\nu\beta}+\tilde{g}_{\mu\beta}\tilde{g}_{\nu\alpha})-\frac{1}{3}\tilde{g}_{\mu\nu}\tilde{g}_{\alpha\beta} \quad (10)$$

for the tensor f_2 meson, where $\tilde{g}_{\mu\nu}=-g_{\mu\nu}+\frac{q^\mu q^\nu}{m^2_{f_2}}$.

As shown in Fig. 2 and 3, we present the variation of the total cross sections compare with the experiment data to the different typical Λ_K values, where Λ_K is taken 0.51 GeV to 0.53 GeV for $K^-p\to f_2(1270)\Lambda$ and $K^-p\to f_2'(1525)\Lambda$. The obtained differential cross section for $K^-p\to f_2(1270)\Lambda$ and $K^-p\to f_2'(1525)\Lambda$ with the typical cutoff Λ_K are also analyzed in Fig 4 and Fig 5, respectively. The experimental data denoted by the squares are taken from Ref. [11]. We see that our theoretical results, which is obtained including the contributions from the $t-$channel K^+ exchange and nucleon pole, can give a reasonable description of the experimental data.

III. THE $K^-p\to f_2(1270)\Lambda, f_2'(1525)\Lambda\to K^+K^-\Lambda$ REACTION

Turning now to the $K^-p\to f_2\Lambda\to K^+K^-\Lambda$ reaction is described Fig. 6, the scattering amplitude $\mathcal{M}_{fi}(K^-p\to K^+K^-\Lambda)$ is

$$\mathcal{M}^a_{fi}(K^-p\to K^+K^-\Lambda)=i(\frac{g_{f_2K^+K^-}}{2m_{f_2}})^2\frac{g_{Kp\Lambda}}{M_N+M_\Lambda}$$
$$\times[p_4^\mu(p_3-p_4)^\nu-(p_3-p_4)^\mu p_3^\nu]G^T_{\mu\nu\alpha\beta}(q)$$
$$\times[k_t^\alpha(p_1-k_t)^\beta-(p_1-k_t)^\alpha p_1^\beta]\frac{1}{k_t^2-m_k^2}$$
$$\times\bar{u}(p_5,s_5)\not{k}_t\gamma_5 u(p_2,s_2)F_K(k_t^2)\mathcal{F}_T(q^2,M_{f_2}), \quad (11)$$
$$\mathcal{M}^b_{fi}(K^-p\to K^+K^-\Lambda)=-i\frac{g_{f_2K^+K^-}}{2m_{f_2}}\frac{g_{Kp\Lambda}}{M_N+M_\Lambda}\frac{g_{f_2NN}}{2M_N}$$
$$\times[p_4^\mu(p_3-p_4)^\nu-(p_3-p_4)^\mu p_3^\nu]G^T_{\mu\nu\alpha\beta}(q)$$
$$\times\bar{u}(p_5,s_5)\not{p}_1\gamma_5\frac{(\not{k}_u+M_N)}{k_\mu^2-M_N^2}(\gamma_\alpha p_{2\beta}+\gamma_\beta p_{2\alpha})$$
$$\times u(p_2,s_2)F_N(k_u^2)\mathcal{F}_T(q^2,M_{f_2}), \quad (12)$$

where $G^T_{\mu\nu\alpha\beta}$ is the propagator of the tensor meson f_2 and can read

$$G^T_{\mu\nu\alpha\beta}=\frac{P_{\mu\nu\alpha\beta}}{q^2-m^2_{f_2}+im_{f_2}\Gamma_{f_2}}. \quad (13)$$



FIG. 6: Feynman diagrams for $K^-p\to f_2(1270)\Lambda, f_2'(1525)\Lambda\to K^+K^-\Lambda$ reaction. In the diagram, we also show the definition of the kinematical (p_1,p_2,p_3,p_4,p_5) that we use in the present calculation.

With the formalism and ingredients given above, the cross section versus the beam momentum P_{K^-} for $K^-p\to K^-K^+\Lambda$ reaction is calculated by using a Monte Carlo multiparticle phase space integration program. Our predictions, with $\Lambda_K=0.51$ to 0.53 GeV, for the beam momentum P_{K^-} from just above the production threshold 1.7 GeV to 6.0 GeV are shown in Fig. 7. The solid and dash curves stand for the contributions from the $f_2(1270)$ and $f_2'(1525)$, respectively.



FIG. 7: (color online). The cross section for $K^-p\to f_2(1270)\Lambda, f_2'(1525)\Lambda\to K^+K^-\Lambda$ reaction.

Furthermore, the corresponding K^+K^- invariant mass spectrum for the $K^-p\to K^+K^-\Lambda$ reaction with $\Lambda_K=0.52$ GeV at beam momentum $P_{K^-}=3.9$ and 4.6 GeV are calculated and shown in Fig 8. The dashed lines are pure phase space distributions, while, the solid lines are full results from our model.



FIG. 8: The K^-K^+ invariant mass spectrum for the $K^-p \to K^+K^-\Lambda$ reaction at beam momentum P_{K^-} = 3.9 and 4.6 GeV. The dashed lines are pure phase space distributions, while, the solid lines are full results from our model.

From Fig 8, we can see that there is a clear peak in the K^-K^+ invariant mass distribution, which is produced by including the contribution from the $f_2(1270)$ and $f_2^{'}(1525)$, respectively.

IV. SUMMARY

In this work, we perform a calculation of the $f_2[\equiv f_2(1270)$ $f_2^{'}(1525)]$ tenser meson production in the $K^-p \to f_2\Lambda$ and $K^-p \to K^-K^+\Lambda$ reaction within the effective Lagrangian method. For the $K^-p \to f_2\Lambda$ reaction, by considering the contributions from the t-channel K^+ exchange and u-channel nucleon pole, we get a fairly good description of the experimental total cross section data. Our model shown the differential cross section $d\sigma/dt$ as function as $-t$, and get a good description of the experimental differential cross section data.

Basing on our results of $K^-p \to f_2\Lambda$ reaction, we have studied the $K^-p \to K^-K^+\Lambda$ reaction. In this case, we have considered the contributions from only the $f_2(1270)$ and $f_2^{'}(1525)$ mesons. The invariant mass distribution for the Dalitz process $K^-p \to K^-K^+\Lambda$ shows an obvious peak at $M^2_{K^+K^-} \approx 1.63$ GeV and $M^2_{K^+K^-} \approx 2.33$ GeV, respectively, which can be checked by further experiment.

Acknowledgments

This project was partially supported by the Major State Basic Research Development Program in China (No. 2014CB845400), the National Natural Science Foundation of China (Grants No. 11475227, No. 11275235, No. 11035006, No.11175220) and the Chinese Academy of Sciences (the Knowledge Innovation Project under Grant No. KJCX2-EW-N01, Century program under Grant No. Y101020BR0). It is also supported by the Open Project Program of State Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Sciences, China (No. Y5KF151CJ1).

[1] R. Aaij *et al.* [LHCb Collaboration], "Observation of J/p Resonances Consistent with Pentaquark States in ^0_b J/K⁻p Decays," Phys. Rev. Lett. **115**, 072001 (2015) doi:10.1103/PhysRevLett.115.072001 [arXiv:1507.03414 [hep-ex]].

[2] K.A. Olive et al. (Particle Data Group), Chin. Phys. C, **38**, 090001 (2014).

[3] L. S. Geng and E. Oset, "Vector meson-vector meson interaction in a hidden gauge unitary approach," Phys. Rev. D **79**, 074009 (2009) doi:10.1103/PhysRevD.79.074009 [arXiv:0812.1199 [hep-ph]].

[4] H. Nagahiro, J. Yamagata-Sekihara, E. Oset, S. Hirenzaki and R. Molina, "The gamma gamma decay of the f(0)(1370) and f(2)(1270) resonances in the hidden gauge formalism," Phys. Rev. D **79**, 114023 (2009) doi:10.1103/PhysRevD.79.114023 [arXiv:0809.3717 [hep-ph]].

[5] A. Martinez Torres, L. S. Geng, L. R. Dai, B. X. Sun, E. Oset and B. S. Zou, "Study of the J/psi → phi(omega) f(2)(1270), J/psi → phi (omega) f-prime(2)(1525) and J/psi → K*0(892) anti-K*0(2) (1430) decays," Phys. Lett. B **680** (2009) 310 doi:10.1016/j.physletb.2009.09.003 [arXiv:0906.2963 [nucl-th]].

[6] L. S. Geng, F. K. Guo, C. Hanhart, R. Molina, E. Oset and B. S. Zou, "Study of the f(2)(1270), f(2)-prime(1525), f(0)(1370) and f(0)(1710) in the J/psi radiative decays," Eur. Phys. J. A **44**, 305 (2010) doi:10.1140/epja/i2010-10971-5 [arXiv:0910.5192 [hep-ph]].

[7] J. J. Xie and E. Oset, "Photoproduction of the $f_2(1270)$ resonance," Eur. Phys. J. A **51**, 111 (2015) doi:10.1140/epja/i2015-15111-3 [arXiv:1412.3234 [nucl-th]].

[8] M. Battaglieri *et al.* [CLAS Collaboration], "Photoproduction of pi+ pi- meson pairs on the proton," Phys. Rev. D **80**, 072005 (2009) doi:10.1103/PhysRevD.80.072005 [arXiv:0907.1021 [hep-ex]].

[9] J. J. Xie, E. Oset and L. S. Geng, "Photoproduction of the $f_2'(1525), a_2(1320)$, and $K_2^*(1430)$," Phys. Rev. C **93**, no. 2, 025202 (2016) doi:10.1103/PhysRevC.93.025202 [arXiv:1509.06469 [nucl-th]].

[10] M. Aguilar-Benitez *et al.* [CERN-College de France-Madrid-Stockholm Collaboration], "Study of Hypercharge Exchange Reactions of the Type 0- 1/2+ → 2+ 1/2+ at 4-GeV/*c*," Z. Phys. C **8**, 313 (1981). doi:10.1007/BF01546327

[11] M. Aguilar-Benitez, S. U. Chung, R. L. Eisner and N. P. Samios, "Study of nonstrange mesons produced in k- p interactions at 3.9 and 4.6 gev/c," Phys. Rev. D **6**, 29 (1972). doi:10.1103/PhysRevD.6.29

[12] J. Mott *et al.*, "Study of k-minus p interactions at 4.1 and 5.5 gev/c - final states with two charged particles and a visible lambda," Phys. Rev. **177**, 1966 (1969). doi:10.1103/PhysRev.177.1966

[13] Y. Oh, C. M. Ko and K. Nakayama, "Nucleon and Delta resonances in K Sigma(1385) photoproduction from nucleons," Phys. Rev. C **77**, 045204 (2008) doi:10.1103/PhysRevC.77.045204 [arXiv:0712.4285 [nucl-th]].

[14] Y. Yan and R. Tegen, "Baryon exchange and meson pole diagrams in p anti-p → anti-K K, pi+ pi-," Nucl. Phys. A **648**, 89 (1999). doi:10.1016/S0375-9474(99)00019-6

[15] H. Goldberg, "Backward (theta=180-degrees) piN Dispersion Relations: Applications to the Interference Model, P and P-prime Trajectories, and to the Mechanical Form Factors of the Nucleon," Phys. Rev. **171**, 1485 (1968). doi:10.1103/PhysRev.171.1485

[16] H. Pilkuhn, W. Schmidt, A. D. Martin, C. Michael, F. Steiner, B. R. Martin, M. M. Nagels and J. J. de Swart, "Compilation of coupling constants and low-energy parameters. 1973 edition," Nucl. Phys. B **65**, 460 (1973). doi:10.1016/0550-3213(73)90296-4

[17] B. Renner, "Empirical test of tensor meson dominance," Phys. Lett. B **33**, 599 (1970). doi:10.1016/0370-2693(70)90359-X

[18] Y. s. Oh and T. S. H. Lee, "rho meson photoproduction at low-energies," Phys. Rev. C **69**, 025201 (2004) doi:10.1103/PhysRevC.69.025201 [nucl-th/0306033].

[19] S. H. Kim, S. i. Nam, Y. Oh and H. C. Kim, "Contribution of higher nucleon resonances to $K^*\Lambda$ photoproduction," Phys. Rev. D **84**, 114023 (2011) doi:10.1103/PhysRevD.84.114023 [arXiv:1110.6515 [hep-ph]].